

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

Final Report Covering October 1973 - April 1975

on

National Aeronautics and Space Administration

Grant NGR-06-004-123

Grant Title: A Study of Geomagnetic Storms

Principal Investigator: Dr. V. L. Patel
Associate Professor
Department of Physics and Astronomy
University of Denver
Denver, Colorado 80210

NASA Technical Officer
for this Grant: Dr. H. D. Calahan, Physics and Astronomy
Code SG, NASA Headquarters

(NASA-CR-142800) A STUDY OF GEOMAGNETIC
STORMS Final Report, Oct. 1973 - Apr. 1975
(Denver Univ.) 4 p HC \$3.25 CSCL 08N

N75-24208

Unclas
G3/46 21790



We have carried out the study of the initial phase of the geomagnetic storm by using interplanetary magnetic field and plasma data from Explorer series satellites. Twenty-one geomagnetic storm events during 1966-70 have been studied by using simultaneous interplanetary magnetic field and plasma parameters. Explorer 33 and 35 field and plasma data have been analyzed on large-scale (hourly) and small-scale (3 min.) during the time interval coincident with the initial phase of the geomagnetic storms. The principal results show that the solar-ecliptic B_z component turns southward at the end of the initial phase, thus triggering the main phase decrease in Dst geomagnetic field. When the B_z is already negative, its value becomes further negative. The B_y component also shows large fluctuations along with B_z . When there are no clear changes in the B_z component, the B_y shows abrupt changes at the main phase onset. On the small-scale behavior of the magnetic field and electric field ($\vec{E} = -\vec{V} \times \vec{B}$) studied in details for the three events, it is found that the field fluctuations in B_y , B_z and E_y and E_z are present in the initial phase. These fluctuations become larger just before the main phase of the storm begins. In the large-scale behavior field remains quiet because the small-scale variations are averaged out. The power spectrum analysis for few events using 5.2 s data show the steepening of the spectrum after the shock wave associated with ssc. It appears that large as well as small time-scale fluctuations in the interplanetary field and plasma help to change the internal electromagnetic state of the magnetosphere so that a ring current could start causing a geomagnetic storm decrease.

Preliminary results were presented at the IAGA assembly in Kyoto, Japan¹ and the final results are published in a technical report of the University of Denver.²

1. V. L. Patel and M. J. Wiskershen, IAGA Bulletin No. 34, 175, 1973.
2. V. L. Patel and M. J. Wiskershen, Univ. of Denv. Dept. of Physics Tech. Report # 7515, 1975.

V-70

FIELD

INITIAL PHASE OF GEOMAGNETIC STORMS AND INTERPLANE-
TARY FIELDS AND PLASMA PARAMETERS

Patel, V. L., Phys. Dept., U. of Denver, Denver, Co., 80210 USA
 Wiskerchen, M. J., Phys. Dept., U. of Denver

Interplanetary magnetic field variations have been studied for 25 geomagnetic storms during a period of 1966-1970. Hourly values of the interplanetary field components in solar-ecliptic coordinates: B_x , B_y , B_z are compared with the hourly Dst index representing storm variations. In most of these events B_z is found to be southward near the beginning of the main phase of the storms. However, there are events in which B_y component becomes more important than B_z component in correlation with Dst during initial and main phases of the storms. We have further examined E_x , E_y and E_z components of the interplanetary electric field $\vec{E} = -\vec{V} \times \vec{B}$ where V is plasma velocity. Detailed analysis of three geomagnetic storms show that both E_y and E_z components have effects on Dst near the main phase. Further analysis using 3 min. averages shows large and rapid fluctuations in the interplanetary electric field components during initial phase of the geomagnetic storms. For these three storm events (Feb. 7, 15-16, 1967; Feb. 10, 1968) detailed analysis of the interplanetary conditions represented by B_x , B_y , B_z , E_x , E_y , E_z and dE/dt for the time intervals corresponding to the initial phase of the storms will be discussed.

PRECEDING PAGE BLANK NOT FILMED

ORIGINAL PAGE IS
 OF POOR QUALITY

ABSTRACT

Twenty-one geomagnetic storm events during 1966-70 have been studied by using simultaneous interplanetary magnetic field and plasma parameters. Explorer 33 and 35 field and plasma data have been analyzed on large-scale (hourly) and small-scale (3 min.) during the time interval coincident with the initial phase of the geomagnetic storms. The solar-ecliptic B_z component turns southward at the end of the initial phase, thus triggering the main phase decrease in Dst geomagnetic field. When the B_z is already negative, its value becomes further negative. The B_y component also shows large fluctuations along with B_z . When there are no clear changes in the B_z component, the B_y shows abrupt changes at the main phase onset. On the small-scale behavior of the magnetic field and electric field ($E = -V \times B$) studied in details for the three events, it is found that the field fluctuations in B_y , B_z and E_y and E_z are present in the initial phase. These fluctuations become larger just before the main phase of the storm begins. In the large-scale behavior field remains quiet because the small-scale variations are averaged out. The power spectrum analysis for few events using 5.2 data show the steepening of the spectrum after the shock wave associated with the passage of ssc. It appears that large as well as small time-scale fluctuations in the interplanetary field and plasma help to alter the internal electromagnetic state of the magnetosphere so that a ring current could start causing a geomagnetic storm decrease.